Poster Walk Poster Walk

Poster of Distinction

Poster Viewing Session 1 Thursday, 12.09.2019 · 12:50 – 13:50 h

Chairs: Raanan Shamir (Israel) • Kaija-Leena Kolho (Finland)

| P-001 | Ju Whi Kim, Republic of Korea |
|-------|---|
| | GM-CSF producing NCR- ILC3s directly activate neutrophils in the intestinal mucosa of inflammatory bowel disease |
| P-002 | Johan Van Limbergen, The Netherlands |
| | Shifts in bacterial community function are associated with Short Chain Fatty Acid (SCFA) pathways during nutritional therapy in paediatric Crohn´s disease patients |
| P-004 | Sara Sila, Croatia |
| | Altered intestinal microbiota is present in newly diagnosed IBD patients and significantly differs from that of healthy siblings and healthy controls |
| P-006 | Eytan Wine, Canada |
| | Increased faecal short chain fatty acid content with Infliximab therapy in paediatric crohn disease |
| P-009 | Corentin Babakissa, Canada |
| | IL-17-related signature genes are linked to human necrotizing enterocolitis |
| P-014 | Ola Olén, Sweden |
| | Paediatric inflammatory bowel diseases and risk of serious infections: A nationwide register-based cohort study |
| P-017 | Ilona Csizmadia, Ireland |
| | Risk factors for developing paediatric inflammatory bowel disease in an Irish prospective cohort |
| P-020 | Tatsuki Mizuochi, Japan |
| | Diagnostic accuracy of serum proteinase 3 antineutrophil cytoplasmic antibodies for paediatric patients with ulcerative colitis: a prospective multicenter study in Japan |
| P-025 | Gili Focht, Israel |
| | The newly simplified MaRIA score (MaRIAs) is as accurate as the original MaRIA in paediatric Crohn´s Disease: a study from the ImageKids cohort |
| P-033 | Valeria Dipasquale, Italy |
| | Pouchitis in pediatric ulcerative colitis: a multicenter study on behalf of Italian society of pediatric gastroenterology, hepatology and nutrition |

Poster Viewing Session 2 Friday, 13.09.2019 · 13:00 – 13:50 h

Chairs: Sanja Kolaček (Croatia) · Sibylle Koletzko (Germany)

| P-047 | Lucia Cococcioni, United Kingdom |
|-------|---|
| | Safety and efficacy of ferric carboxymaltose [FCM] for the treatment of iron deficiency anaemia in paediatric patients affected by inflammatory bowel disease (pIBD) |
| P-052 | Thomas Walters, Canada |
| | Exclusive enteral nutrition versus corticosteroid induction therapy for new onset paediatric Crohn's disease: comparison of 18 month outcomes in a Canadian prospective multi-centre inception cohort |
| P-066 | Ryma Iratni, Canada |
| | Trends in TNF-alpha inhibitor utilization in children with IBD during the last 10 years: 2009–2018 |
| P-068 | Maria Myrthe Elisabeth Jongsma, The Netherlands |
| | Infliximab in young pediatric IBD patients: it is all about the dosing – a multicenter study from the pediatric IBD Porto group of ESPGHAN |
| P-071 | Ben Kang, Republic of Korea |
| | Infliximab trough levels are not associated with transmural healing at 1 year treatment in paediatric Crohn's disease patients |
| P-076 | Alexandra Petrova, Canada |
| | Infliximab clearance at and post induction predicts clinical and endoscopic outcomes at 1 year in children with Crohn's disease treated with Infliximab |
| P-103 | Patrick van Rheenen, The Netherlands |
| | Time-to-reach target calprotectin level and relation with first relapse in newly diagnosed patients with IBD: first results of the Fast Forward Care Prospective Registry |
| P-110 | Martha. A.C van Gaalen, The Netherlands |
| | Transition test is a valid tool to monitor disease knowledge in adolescents with IBD transitioning to adult care |

5th International Symposium on PIBD www.pibdcongress.org